arXiv:1506.09145v1 [math.CO] 30 Jun 2015

Track Layouts, Layered Path Decompositions, and Leveled Planarity*

Michael J. Bannister[1], William E. Devanny[1], Vida Dujmović[2], David Eppstein[1], and David R. Wood[3]

[1] Department of Computer Science, University of California, Irvine
Irvine, California, USA, {mbannist,wdevanny,eppstein}@uci.edu
[2] School of Computer Science and Electrical Engineering, University of Ottawa
Ottawa, Canada, vida.dujmovic@uottawa.ca
[3] School of Mathematical Sciences, Monash University
Melbourne, Australia, david.wood@monash.edu

Abstract. We investigate two types of graph layouts, track layouts and layered path decompositions, and the relations between their associated parameters track-number and layered pathwidth. We use these two types of layouts to characterize leveled planar graphs, the graphs with planar layered drawings with no dummy vertices. It follows from the known NP-completeness of leveled planarity that track-number and layered pathwidth are also NP-complete, even for the smallest constant parameter values that make these parameters nontrivial. We prove that the graphs with bounded layered pathwidth include outerplanar graphs, Halin graphs, and squaregraphs, but that (despite having bounded track-number) series-parallel graphs do not have bounded layered pathwidth. Finally, we investigate the parameterized complexity of these layouts, showing that past methods used for book layouts don't work to parameterize the problem by treewidth or almost-tree number but that the problem is (non-uniformly) fixed-parameter tractable for tree-depth.

1 Introduction

A *k-track layout* of a graph is a partition of the vertices into k ordered independent sets called *tracks*, and a partition of the edges into non-crossing subsets that connect pairs of tracks. The *track-number* of a graph is the minimum k for which it has a k-track layout. Track layouts are connected with the existence of low-volume three-dimensional graph drawings: a graph has a three-dimensional drawing in an $O(1) \times O(1) \times O(n)$ grid if and only if it has track-number $O(1)$ [6,10]. In this paper we show that track layouts are also related to a more abstract structure in graphs, a *layered path decomposition*. This is a path decomposition together with a partition of the vertices of the graph into a sequence of layers,

* Michael J. Bannister and David Eppstein were supported in part by NSF grant CCF-1228639. William E. Devanny was supported by an NSF Graduate Research Fellowship under grant DGE-1321846. Vida Dujmović was supported by NSERC. David Wood was supported by the Australian Research Council.

where the endpoints of each edge belong to a single layer or two consecutive layers. The *width* of a layered path decomposition is the size of the largest intersection between a bag of the decomposition and a layer. The *layered pathwidth* of a graph is the minimum width of a layered path decomposition.

Our results include the following:

- Both track layouts and layered path decompositions can be used to characterize *leveled planar graphs*, undirected graphs that can be given a Sugiyama-style layered graph drawing with no crossings and no dummy vertices. Specifically, we prove that the leveled planar graphs are exactly the graphs with layered pathwidth at most 1, and are exactly the bipartite graphs with track-number at most 3. Based on the known NP-completeness of testing leveled planarity [17], it follows that testing whether the track-number is at most 3 is NP-complete. This solves an open problem from 2004 [9]. In addition, it implies that testing whether the layered pathwidth is at most 1 is also NP-complete.
- We show that bipartite outerplanar graphs and squaregraphs have layered pathwidth 1, and arbitrary outerplanar graphs and Halin graphs have layered pathwidth ≤ 2. On the other hand, series-parallel graphs (and even the tree-apex graphs, a subclass of series-parallel graphs formed by adding a single vertex to a tree) do not have bounded layered pathwidth, even though they do have bounded track-number.
- We show that, more generally, graphs of bounded layered pathwidth always have bounded track-number. For track-number at most 3, we conjecture that the reverse is true, contrasting the fact that there exist graphs of track-number 4 and unbounded layered pathwidth.
- We show that known methods of obtaining fixed-parameter tractable algorithms for other types of planar embedding, based on Courcelle's Theorem for treewidth [2], or on kernelization of the 2-core for k-almost-trees [3], do not generalize to leveled planarity, track-number, or layered pathwidth. However, for any fixed bound on the tree-depth of an input graph, these problems can be solved in linear time.

2 Definitions

A *track layout* of a graph is a partition of its vertices into sequences, called *tracks*, such that the vertices in each sequence form an independent set and the edges between each two pairs of tracks form a non-crossing set. This means that there do not exist edges uv and $u'v'$ such that u is before u' in one track, but v is after v' in another track; such a pair of edges is said to form a *crossing*. (This ordering constraint on endpoints of pairs of edges connecting two tracks is the same as the constraint on the left-to-right ordering within levels on the endpoints of two edges connecting the same two levels of a layered drawing.)

The *track-number* of a graph G is the minimum number of tracks in a track layout of G; this is finite, since the layout in which each vertex forms its own track is always non-crossing. The set of edges between two tracks form a forest

of caterpillars (a forest in which the non-leaf vertices of each component induce a path); in particular, the graphs with track-number 1 are the independent sets, and the graphs with track-number 2 are the forests of caterpillars [15].

A *tree-decomposition* of a graph G is given by a tree T whose nodes index a collection $(B_x \subseteq V(G) : x \in V(T))$ of sets of vertices in G called *bags*, such that:

- For every edge vw of G, some bag B_x contains both v and w, and
- For every vertex v of G, the set $\{x \in V(T) : v \in B_x\}$ induces a non-empty (connected) subtree of T.

The *width* of a tree-decomposition is $\max_x |B_x| - 1$, and the *treewidth* of a graph G is the minimum width of any tree decomposition of G. Treewidth was introduced (with a different but equivalent definition) by Halin [14] and tree decompositions were introduced by Robertson and Seymour [19].

A *layering* of a graph is a partition of the vertices into a sequence of disjoint subsets (called *layers*) such that each edge connects vertices in the same layer or consecutive layers. One way, but not the only way, to obtain a layering is the *breadth first layering* in which we partition the vertices by their distances from a fixed starting vertex, using breadth-first search [7,8].

A *layered tree decomposition* of a graph is a tree decomposition together with a layering. The *layered width* of layered tree decomposition is the size of the largest intersection of a bag with a layer. The *layered treewidth* of a graph G is the minimum layered width of a tree-decomposition of G. Dujmović, Morin, and Wood [7,8] introduced layered treewidth and proved amongst other results that every planar graph has layered treewidth at most 3, and more generally, that every graph with Euler genus g has layered treewidth at most $2g + 3$.

A *path decomposition* is a tree decomposition where the underlying tree is a path [20]. Thus, it can be thought of as a sequence of subsets of vertices (called *bags*) such that each vertex belongs to a contiguous subsequence of bags and each two adjacent vertices have at least one bag in common. *Layered path decomposition* and *layered pathwidth* are defined in an analogous way to layered tree decomposition and layered treewidth. So the *layered pathwidth* of a graph G is the minimum integer k such that for some path decomposition and layering of G, the intersection of each bag with each layer has at most k vertices. The present paper is the first to consider layered path decompositions.

3 Leveled planarity

The class of *leveled planar graphs* was introduced in 1992 by Heath and Rosenberg [17] in their study of queue layouts of graphs. A leveled planar drawing of a graph is a planar drawing in which the vertices are placed on a collection of parallel lines, and each edge must connect vertices in two consecutive parallel lines. Another equivalent way to state this is that this kind of drawing is a Sugiyama-style layered drawing [22] that achieves perfect quality according to two of the most important quality measures for the drawing, the number of edge crossings [12] and the number of dummy vertices [16].




Fig. 1. Converting a layered drawing to a 3-track layout

In Appendix A we show that the graphs that automatically have leveled planar drawings include several natural and well-studied classes of graphs, including the bipartite outerplanar graphs, squaregraphs, and dual graphs of arrangements of monotone curves. As we will show, the graphs with leveled planar drawings can be characterized both by their low track-number and their low layered pathwidth. This, together with the fact that recognizing leveled planar graphs is NP-complete [17], will imply that testing whether the track-number or layered pathwidth is small is also NP-complete.

3.1 Track layouts

Lemma 1 (implicit in [13]). *Every leveled planar graph has a 3-track layout.*

Proof. Assign the vertices of the graph to tracks according to the number of their level in the layered drawing, modulo 3, as shown in Fig. 1. Within each track, order the vertices within each level contiguously, and order the levels by their positions in the layered drawing. Two edges that connect the same pair of levels cannot cross because of the chosen vertex ordering within the levels, and two edges that connect different pairs of levels but are mapped to the same pair of tracks cannot cross because of the ordering of the levels within the tracks. □

Lemma 1 can be interpreted as 'wrapping' a layered drawing on to 3 tracks; see [9] for a more general wrapping lemma. As Fig. 1 shows, a 3-track layout can also be interpreted geometrically, as a planar drawing in which the tracks are represented as three rays from the origin; it follows from this interpretation that 3-track graphs (and the weakly leveled planar graphs described in Section 4) have universal point sets of size $O(n)$, consisting of n points on each ray. However, for more than three tracks, a similar embedding of the tracks as rays in the plane would not lead to a planar drawing, because there is no requirement that edges of the graph connect only consecutive rays. Indeed, all graphs (for example, arbitrarily large complete graphs) have 4-track subdivisions [11].

Define an *arc* of an undirected graph G to be a directed edge formed by orienting one of the edges of G. For a graph G with a 3-track layout, define a function δ from arcs to ± 1 as follows: if an arc uv goes from track i to track $i+1 \pmod 3$ (that is, if it is oriented clockwise in the planar embedding described above), let $\delta(uv) = +1$; otherwise (if it is oriented counterclockwise), let $\delta(uv) = -1$. For an oriented cycle C, we define (by abuse of notation) $\delta(C) = \sum_{uv \in C} \delta(uv)$.

Lemma 2. *Let C be a cycle embedded in a 3-track layout. Cyclically orient the edges of C. If C is even then $\delta(C) = 0$. If C is odd then $|\delta(C)| = 3$.*

Proof. We proceed by induction on $|C| := |V(C)|$. If $|C| = 3$, then C has one vertex on each track and $\delta(C) \in \{3, -3\}$. If $|C| = 4$, then C has two edges with $\delta = +1$ and two edges with $\delta = -1$, implying $\delta(C) = 0$. Now assume that $|C| \geq 5$. Use the 3-track layout to embed C in the plane as described above, but with straight edges instead of the curved edges shown in the figure. As a planar polygon, C has at least two ears, triangles formed by two of its edges that are empty of other vertices of C (which may be found as the leaf edges in the tree formed as the dual graph of a triangulation of C). If one ear has the same sign of δ for both of the edges that form it, these edges must connect pairs of vertices that are the innermost on their tracks. Therefore, two such ears with same-sign edges could only exist if C is a triangle. For any longer cycle, let uvw be an ear for which $\delta(uv) = -\delta(vw)$; thus edges uv and vw both connect the same two tracks, and (by the assumption that triangle uvw is empty) u and w are consecutive in their track. By deleting v and merging uw into a single vertex, we construct a cycle C' with $|C'| = |C| - 2$, and a 3-track layout of C' with $\delta(C') = \delta(C)$. The result follows by induction. □

The previous lemma is akin to showing that every even oriented cycle C has the winding number 0 and every odd cycle the winding number 1 around the origin of a 3-track representation of C with three rays, Fig. 1.

While Lemma 1 shows that a leveled planar drawing can be wrapped on to three tracks, we now use Lemma 2 to show that a bipartite 3-track layout can be unwrapped to produce a leveled planar drawing.

Theorem 1. *A graph G has a leveled planar drawing if and only if G is bipartite and has a 3-track layout.*

Proof. In one direction, if G has a leveled planar drawing, then it is bipartite (with a coloring determined by the parity of the level numbers of the drawing) and has a 3-track layout by Lemma 1.

In the other direction, suppose that G is bipartite and has a 3-track layout. We may assume without loss of generality that G is connected, for otherwise we can draw each connected component of G separately; let T be a spanning tree of G, and let v be an arbitrary vertex of G. Assign v to level zero of a layered drawing, and assign each other vertex w to the level given by the sum of the numbers $\delta(xy)$ for the edges xy of the oriented path from v to w in T. (Some of these level numbers may be negative.) By construction, the endpoints of each

edge of T are assigned to consecutive levels, and by applying Lemma 2 to the oriented cycle formed by a non-tree edge together with the tree path connecting its endpoints, the same can be shown to be true of each edge of $G - E(T)$.

Within each level of the drawing, the vertices all come from the same track, determined by the value of the level modulo 3. Assign the vertices to positions in left-to-right order on this level according to their ordering within this track. Then no two consecutive levels of the drawing can have crossing edges, because such a crossing would also be a crossing in the track layout. Therefore, this assignment of vertices to levels and to positions within these levels gives a leveled planar drawing of G. □

3.2 Layered pathwidth

The following lemma will allow us to build a layered path decomposition of a leveled planar graph greedily, one bag at a time.

Lemma 3. *Let G be a graph with a leveled planar drawing, and let S be a subset of vertices of G containing one vertex s_i in each layer i of the drawing. Suppose also that there exists at least one vertex in S that is not the rightmost vertex in its layer. Then there exists i such that s_i is not rightmost in its layer, and such that each neighbor of s_i either belongs to S or is positioned to the left of a vertex in S within its layer.*

Proof. If every vertex in S is either rightmost in its level, or has no neighbor to the right of S, then we are done, for we may choose s_i to be any vertex that is not rightmost in its level.

Otherwise, draw a directed graph D whose vertices are the layers of G, with an edge from layer i to layer j ($j = i \pm 1$) if s_i has a neighbor in layer j to the right of s_j. Then D cannot contain edges in both directions between s_i and s_j, for the corresponding edges in G would necessarily cross, so D must be a subgraph of an oriented path, and in particular must be a directed acyclic graph. By the assumption that at least one vertex has a neighbor to the right of S, D has at least one edge. Therefore, D contains a vertex i (a level of the drawing) that has incoming edges but that does not have any outgoing edges. For this level, s_i is not rightmost in its level (else i could have no incoming edges) but has no neighbors to the right of S (else i would have an outgoing edge), as desired. □

Theorem 2. *A graph G is leveled planar if and only if it has layered pathwidth 1.*

Proof. In one direction, suppose that G has layered pathwidth 1. We can construct a layered drawing of G from its layered path decomposition, by using the layers of the layered path decomposition as the levels of the layered drawing. Because the pathwidth is 1, any two vertices in the same level occur within disjoint intervals of the sequence of bags of the path decomposition of G, and so we can order the vertices within each level of the drawing by the ordering of their bags in the path decomposition. Each edge of G must connect two consecutive levels of this drawing (no edge can connect two edges in the same level because then

the bag containing its endpoints would intersect that level in a set of size two or more), and the edges can be ordered from left to right by the ordering of the bags containing their endpoints. This ordering is consistent with the ordering of the endpoints in both levels spanned by the edge, so there can be no crossings.

In the other direction (implicit in [5, Lemma 1]), suppose that G has a leveled planar drawing. We must show that this information can be used to find a layered path-decomposition of G with width 1. For the layering of this layered path-decomposition, we use the sequence of levels of the drawing of G; because the drawing is assumed to have no dummy vertices, this satisfies the definitional requirement of a layering, that each edge connect vertices in the same layer or in two consecutive layers. For the path decomposition, we use a sequence of bags with one vertex per layer, this first of which is the set of vertices that are leftmost in their layer of the drawing. We construct this sequence of bags using a greedy algorithm from this starting bag, at each step using Lemma 3 to find a vertex v whose neighbors all belong to the present bag or earlier bags, and forming the next bag by replacing v with the next vertex to the right of v in the same level. In this way, by construction, each vertex belongs to a consecutive subsequence of bags. Each edge vw has both neighbors in at least one bag (the first bag in the sequence to include the second of its two endpoints), because until the second endpoint has been introduced as part of the sequence of bags, the first endpoint cannot be replaced. Thus, we have a path decomposition with one vertex per layer, showing that the layered pathwidth is 1. □

We conjecture that every 3-track graph (without restriction on bipartiteness) has bounded layered pathwidth.

4 Weakly leveled planarity

We define a *weakly leveled planar drawing* of a graph to be a planar drawing in which the vertices are placed on a sequence of parallel lines and each edge connects two vertices that either belong to the same line or are on consecutive lines. I.e., we relax the definition of leveled planar drawings to allow edges between consecutive vertices on the same level as each other.

Theorem 3. *If a graph G has a weakly leveled planar drawing, then G has layered pathwidth at most 2.*

Proof. The proof is almost the same as that of Theorem 2. We use the sequence of levels of the drawing as the layers of a layered path-decomposition, and form a sequence of bags with one vertex per layer, covering the edges that connect two consecutive layers. As in Theorem 2 we construct this sequence of bags greedily, using Lemma 3 to find a vertex v whose neighbors on adjacent levels all belong to the present bag or earlier bags. In the proof of Theorem 2, the next bag was formed by replacing v with the next vertex w to the right of v on the same level, but if we did that we might form a sequence of bags that did not include a bag containing both vw, violating the definition of a path decomposition if v and w

are adjacent. Instead, we first add w to the present bag, forming a bag whose intersection with v's layer has two vertices, and then we form a second bag by removing v. The result is a path decomposition: every edge between consecutive levels is represented in at least one bag by the same reasoning as in the proof of Theorem 2, and every edge with two endpoints on the same level is represented in at least one bag by construction. Its largest intersection with a level has size two, so the layered pathwidth is 2. □

As a corollary of this theorem together with Lemma 4 (below), weakly leveled planar graphs also have bounded track-number. In Appendix A we show that the graphs with weakly leveled planar drawings include the outerplanar graphs and the Halin graphs, but not the series-parallel graphs.

5 Beyond leveled planarity

Dujmović, Morin, and Wood [6] proved that every graph with pathwidth k has track-number at most $k+1$. Here we provide the following generalisation.

Lemma 4. *Every graph with layered pathwidth ℓ has track-number at most 3ℓ.*

Proof. Let $B_1, B_2, \ldots, B_n$ be a path decomposition of G with layered width ℓ. Let $(V_0, V_1, \ldots, V_m)$ be the corresponding layering. Thus, each bag B_i contains at most ℓ vertices in each layer V_j. Since $G[V_j]$ has pathwidth at most $\ell - 1$, there is a proper colouring of $G[V_j]$ with colours $1, 2, \ldots, \ell$. For each vertex v of G, let $b(v) := \min\{i : v \in B_i\}$ be the index of the leftmost bag containing v. For $0 \leq j \leq m$ and $1 \leq a \leq \ell$, let $V_{j,a}$ be the set of vertices in V_j coloured a. Let $\preceq$ be the total order of $V_{j,a}$ defined by $v \prec w$ if and only if $b(v) < b(w)$. Clearly $\preceq$ is a total order. Since $G[V_j]$ is properly coloured, $V_{j,a}$ is a track. Suppose on the contrary that vw and xy form an X-crossing between $V_{j,a}$ and $V_{k,b}$, where $v \prec x$ in $V_{j,a}$ and $y \prec w$ in $V_{k,b}$. Without loss of generality, $b(w) \leq b(x)$. Since $y \prec w$ we have $b(y) < b(w) \leq b(x)$. Since xy is an edge, $y \in B_{b(w)}$. Hence y and w are adjacent in $G[V_k]$, which is a contradiction since y and w are assigned the same colour. Therefore there is no X-crossing, and $\{V_{j,a} : 0 \leq j \leq m, 1 \leq a \leq \ell\}$ is a track layout of G. Since $(V_0, V_1, \ldots, V_m)$ is a layering, if vw is an edge of G with $v \in V_{j,a}$ and $w \in V_{k,b}$, then $|j - k| \leq 1$. It follows from a result of Dujmović, Por and Wood [9, Lemma 6 with $s = 1$] that this track layout can be wrapped onto 3ℓ tracks. In particular, for $q \in \{0, 1, 2\}$ and $1 \leq a \leq \ell$, let $W_{q,a}$ be the track $V_{q,a}, V_{q+3,a}, V_{q+6,a}, \ldots$. Then $\{W_{q,a} : q \in \{0, 1, 2\}, 1 \leq a \leq \ell\}$ is a 3ℓ-track layout of G. □

There is a natural connection between layered treewidth and layered pathwidth.

Lemma 5. *Every n-vertex graph G with layered treewidth ℓ has layered pathwidth at most $\ell \log_3(2n+1)$.*

Proof. Let $(B_x : x \in V(T))$ be a tree decomposition of G with layered width ℓ. That is, each bag B_x contains at most ℓ vertices in some layering. If $B_x = B_y$ for some edge $xy \in E(T)$, then contracting xy gives a tree decomposition with layered width ℓ. Thus, we may assume that $B_x \neq B_y$ for each edge $xy \in E(T)$. It follows that T has at most n vertices. Scheffler [21] proved that every n-vertex tree has pathwidth at most $\log_3(2n+1)$. Let $B_1, \dots, B_m$ be a path decomposition of T with width $\log_3(2n+1)$. Let $B'_i := \cup\{V_x : x \in B_i\}$. Then $B'_1, \dots, B'_m$ is a path decomposition of G with layered width at most $\ell \log_3(2n+1)$ (with respect to the initial layering). □

Lemmas 4 and 5 imply the following result, which improves the constant factor in a result of Dujmović [4].

Theorem 4. *Every n-vertex graph with layered treewidth ℓ has track-number at most $3\ell \log_3(2n+1)$.*

Note that, similar to layered treewidth [8], layered pathwidth is also not a minor-closed parameter. For example, it is easily seen that the $n \times n \times 2$ grid graph has layered pathwidth at most 3, but G contains a K_n minor [23], and K_n has layered pathwidth $\lceil n/2 \rceil$.

6 Parameterized complexity

A fixed-parameter tractable problem is also *strongly uniform fixed-parameter tractable*. A problem is *uniformly fixed-parameter tractable* if there is an algorithm that solves it in polynomial time for any value of the parameter, but we cannot compute the dependence on the parameter. Lastly a problem is *non-uniformly fixed-parameter tractable* if there is a collection of algorithms such that for each possible value of the parameter one of the algorithms solves the problem in polynomial time.

6.1 Treewidth

We sketch an argument as to why it is not possible to use Courcelle's Theorem (or any automata methods based on tree decompositions) to produce a fixed-parameter tractable algorithm for leveled planarity with respect to treewidth. Consider the family of graphs depicted in Fig. 2. These graphs have bounded treewidth (in fact pathwidth at most 12) and are leveled planar precisely when p equals q. However, since p and q are unbounded it is necessary to carry more than a finite amount of state between bags in a treewidth decomposition when parsing the decomposition. Thus, the decompositions corresponding to leveled planar graphs cannot be recognized by automata or methods using automata such as Courcelle's Theorem. This intuitive argument is made formal using the Myhill-Nerode Theorem for tree automata in Appendix B.




Fig. 2. A family of graphs with bounded treewidth demonstrating that the family of leveled planar graphs is not finite state.




Fig. 3. A family of 2-almost trees for which the standard kernelization cannot decide leveled planarity. The subgraphs T_ℓ are complete binary trees of depth ℓ.

6.2 Tree-depth

The *tree-depth* of a graph G is the minimum height of a forest of rooted trees on the same vertex set as G such that edges in G only go from ancestors to descendants in the forest. It is bounded by pathwidth, and therefore by track-number: track-number$(G) \leq$ pathwidth$(G) + 1 \leq$ tree-depth(G); see [6,18].

Theorem 5. *Computing the track-number or layered pathwidth of a graph G is non-uniformly fixed-parameter linear in the tree-depth of G.*

Proof. Track-number and layered pathwidth are both monotone (cannot increase) under taking induced subgraphs. The graphs with tree-depth bounded by a constant are well-quasi-ordered under taking induced subgraphs and so for any fixed bound on tree-depth and either track-number or layered pathwidth there exist only a finite number of forbidden induced subgraphs [18]. Since the track-number and pathwidth are both bounded by the tree-depth, the same is true for any fixed bound on tree-depth, regardless of track-number or layered pathwidth.

Because induced subgraph testing is linear time for graphs with tree-depth bounded by a fixed number d, we can for each $t \leq d$ test if the graph has any of the forbidden induced subgraphs to track-number t each in linear time [18]. □

However, this argument does not tell us how to find the set of forbidden minors, nor what the dependence of the time bound on the tree-depth is. It would be of interest to replace this existence proof with a more constructive algorithm.

6.3 Almost-trees

The *cyclomatic number* (also called *circuit rank*) of a graph is defined to be $r = m - n + c$ where c is the number of connected components in the graph.

We say that a graph G is a *k-almost tree* if every biconnected component of G has cyclomatic number at most k. The problems of 1-page and 2-page crossing minimization and testing 1-planarity were shown to be fixed-parameter tractable with respect to the k-almost tree parameter, via the kernelization method [1,3].

In these previous papers, the "standard kernelization" used for a k-almost tree G is constructed by first iteratively removing vertices of degree one vertices until no more remain, leaving what is called the *2-core* of G. The 2-core consists of vertices of degree greater than two and paths of degree two vertices connecting these high degree vertices. The paths of degree two vertices are then shortened to a maximum length whose value is a function of k, with a precise form that depends on the specific problem.

However, this kernelization cannot be used to produce a fixed-parameter tractable algorithm for deciding leveled planarity. To see this, consider the graph in Fig. 3, constructed by drawing $K_{2,3}$ in the plane, and replacing each of the three vertices with paths of k vertices, and then rooting a complete binary tree of depth ℓ at one of the vertices of each of these paths. We note that, as complete binary trees have unbounded pathwidth, they also require an unbounded number of layers (depending on ℓ) in any leveled planar drawing. Additionally, depending on the planar embedding chosen for this graph, at most two of its three trees can be drawn on the outside face. So this graph is leveled planar precisely when ℓ is small enough for the remaining tree T_ℓ to be drawn within one of the four bounded faces of the drawing, i.e., the leveled planarity of of the graph depends on the relationship between k and ℓ. Since this relationship is not preserved in the kernelization it can not be used to produce a fixed-parameter tractable algorithm for leveled planarity.

References

1. Michael J. Bannister, Sergio Cabello, and David Eppstein, *Parameterized complexity of 1-planarity*, Algorithms and Data Structures, LNCS, vol. 8037, Springer, 2013, pp. 97–108, doi:10.1007/978-3-642-40104-6_9.
2. Michael J. Bannister and David Eppstein, *Crossing minimization for 1-page and 2-page drawings of graphs with bounded treewidth*, 22nd Int. Symp. Graph Drawing (GD 2014), LNCS, vol. 8871, Springer-Verlag, 2014, pp. 210–221, doi:10.1007/978-3-662-45803-7_18.
3. Michael J. Bannister, David Eppstein, and Joseph A. Simons, *Fixed parameter tractability of crossing minimization of almost-trees*, 21st Int. Symp. Graph Drawing (GD 2013), LNCS, vol. 8242, Springer-Verlag, 2013, pp. 340–351, doi:10.1007/978-3-319-03841-4_30, MR 3162035.
4. Vida Dujmović, *Graph layouts via layered separators*, J. Combin. Theory Series B. **110** (2015), 79–89, doi:10.1016/j.jctb.2014.07.005, MR 3279388.
5. Vida Dujmović, Michael R. Fellows, Matthew Kitching, Giuseppe Liotta, Catherine McCartin, Naomi Nishimura, Prabhakar Ragde, Frances Rosamond, Sue Whitesides, and David R. Wood, *On the parameterized complexity of layered graph drawing*, Algorithmica **52** (2008), no. 2, 267–292, doi:10.1007/s00453-007-9151-1, MR 2434968.

6. Vida Dujmović, Pat Morin, and David R. Wood, *Layout of graphs with bounded tree-width*, SIAM J. Comput. **34** (2005), no. 3, 553–579, doi:10.1137/S0097539702416141, MR 2137079.
7. ———, *Layered separators for queue layouts, 3D graph drawing and nonrepetitive coloring*, Proc. 54th Symp. on Foundations of Computer Science (FOCS 2013), IEEE Computer Society, 2013, pp. 280–289, doi:10.1109/FOCS.2013.38.
8. ———, *Layered separators in minor-closed families with applications*, Electronic preprint arXiv:1306.1595, 2013.
9. Vida Dujmović, Attila Pór, and David R. Wood, *Track layouts of graphs*, Discrete Math. Theor. Comput. Sci. **6** (2004), no. 2, 497–521, available from http://www.dmtcs.org/pdfpapers/dm060221.pdf, MR 2180055.
10. Vida Dujmović and David R. Wood, *Three-dimensional grid drawings with sub-quadratic volume*, Towards a theory of geometric graphs (János Pach, ed.), Contemp. Math., vol. 342, Amer. Math. Soc., 2004, pp. 55–66, doi:10.1090/conm/342/06130, MR 2065252.
11. Vida Dujmović and David R. Wood, *Stacks, queues and tracks: layouts of graph subdivisions*, Discrete Math. Theor. Comput. Sci. **7** (2005), no. 1, 155–201, available from http://www.dmtcs.org/pdfpapers/dm070111.pdf, MR 2164064.
12. Peter Eades and Nicholas C. Wormald, *Edge crossings in drawings of bipartite graphs*, Algorithmica **11** (1994), no. 4, 379–403, doi:10.1007/BF01187020.
13. Stefan Felsner, Giussepe Liotta, and Stephen K. Wismath, *Straight-line drawings on restricted integer grids in two and three dimensions*, J. Graph Algorithms Appl. **7** (2003), no. 4, 363–398, doi:10.7155/jgaa.00075.
14. Rudolf Halin, *S-functions for graphs*, J. Geometry **8** (1976), no. 1-2, 171–186, doi:10.1007/BF01917434, MR 0444522.
15. Frank Harary and Allen Schwenk, *A new crossing number for bipartite graphs*, Utilitas Math. **1** (1972), 203–209.
16. Patrick Healy and Nikola S. Nikolov, *How to layer a directed acyclic graph*, 9th Int. Symp. Graph Drawing (GD 2001), LNCS, vol. 2265, Springer-Verlag, 2002, pp. 16–30, doi:10.1007/3-540-45848-4_2, MR 1962416.
17. Lenwood S. Heath and Arnold L. Rosenberg, *Laying out graphs using queues*, SIAM J. Comput. **21** (1992), no. 5, 927–958, doi:10.1137/0221055, MR 1181408.
18. Jaroslav Nešetřil and P Ossona de Mendez, *Sparsity (graphs, structures, and algorithms)*, Algorithms and Combinatorics, vol. 28, Springer, 2012, doi:10.1007/978-3-642-27875-4.
19. Neil Robertson and P. D. Seymour, *Graph minors. II. Algorithmic aspects of tree-width*, J. Algorithms **7** (1986), no. 3, 309–322, doi:10.1016/0196-6774(86)90023-4, MR 855559.
20. Neil Robertson and Paul Seymour, *Graph minors, I: Excluding a forest*, J. Combinatorial Theory, Ser. B **35** (1983), no. 1, 39–61, doi:10.1016/0095-8956(83)90079-5.
21. Petra Scheffler, *Optimal embedding of a tree into an interval graph in linear time*, Fourth Czechoslovakian Symposium on Combinatorics, Graphs and Complexity, Annals of Discrete Mathematics, vol. 51, Elsevier, 1992, pp. 287–291, doi:10.1016/S0167-5060(08)70644-7.
22. Kozo Sugiyama, Shôjirô Tagawa, and Mitsuhiko Toda, *Methods for visual understanding of hierarchical system structures*, IEEE Trans. Systems, Man, and Cybernetics **SMC-11** (1981), no. 2, 109–125, doi:10.1109/TSMC.1981.4308636, MR 0611436.
23. David R. Wood, *Clique minors in cartesian products of graphs*, New York J. Math. **17** (2011), 627—682.




Fig. 4. Examples of graphs with planar breadth-first layerings (start vertex shown in red, and layering in yellow): left, a bipartite outerplanar graph (Theorem 6); center, a squaregraph (Theorem 7); and right, the dual graph of an arrangement of doubly-unbounded monotonic curves (Theorem 8).

A Special classes of graphs

We consider here particular graph families such as the outerplanar graphs, and prove that these families are leveled planar or weakly leveled planar. Our results are based on breadth-first layerings; we define a layering of a graph to be *planar* if there exists a non-crossing layered drawing of the graph in which the layers of the drawing are the same as the layers of the layering.

A.1 Bipartite outerplanar graphs

Theorem 6 (implicit in [13]). *Every bipartite outerplanar graph is leveled planar. Every breadth first layering of such a graph G gives a leveled planar drawing.*

Proof. Let v be the starting vertex of a breadth first layering. Then for each face cycle C of the outerplanar embedding of G, there must be a unique nearest neighbor in C to v. For, if v were nearest to distinct vertices u and w in C, then by bipartiteness these two vertices must be non-adjacent in C. In this case, the graph formed by C together with the shortest paths from v to u and w would contain a subdivision of $K_{2,3}$ (with u and w as the degree three vertices, two paths between them in C, and one more path between them through the shortest path tree rooted at v), an impossibility for an outerplanar graph. For the same reason, the distances in v from this nearest neighbor or pair of nearest neighbors must increase monotonically in both directions around C until reaching a unique farthest neighbor, because in the same way any non-monotonicity could be used to construct a subdivision of $K_{2,3}$.

Thus, each face cycle of G has a planar breadth first layering. The result follows from the fact that in a plane graph with an assignment of levels to the

vertices, there is a planar drawing consistent with this level assignment and with the given embedding of the graph, if and only if every face cycle of the given graph has a planar drawing consistent with the level assignment [24]. □

A.2 Squaregraphs

A *squaregraph* is defined to be a graph that has a planar embedding in which each bounded face is a 4-cycle and each vertex either belongs to the unbounded face or has four or more incident edges. These graphs may also be characterized in various other ways, for instance as the dual graphs of hyperbolic line arrangements with no three mutually-intersecting lines [25].

Theorem 7. *Every squaregraph G is leveled planar, with a leveled planar drawing coming from a breadth first layering.*

Proof. Because all their bounded faces are even-sided, squaregraphs are necessarily bipartite, so every choice of starting vertex gives a valid breadth first layering. Bandelt et al [25, Lemma 12.2] prove that, for every choice of starting vertex, we can add extra edges to the squaregraph to form a plane multigraph in which the added edges link each layer into a cycle, and in which these cycles are all nested within each other.

Now, choose the starting vertex v to be a vertex of the outer face. Then each cycle added in this augmentation of G contains an edge that separates v from the unbounded face of the augmented graph. If we remove each such edge from the augmented graph, we break each cycle into a path in a consistent way, such that the path ordering within each layer matches the given planar embedding of G. □

A.3 Dual graphs of monotone curves

Theorem 8. *Let A be a collection of finitely many x-monotone curves in the plane, whose projections onto the x-axis cover the entire axis, such that any two curves intersect at finitely many crossing points. Then the dual graph of the arrangement of the curves in A has a planar breadth first layering.*

Proof. Each vertex of the dual graph corresponds to a connected component of the complement of $\bigcup A$; we call this the *region* of the vertex. We may assign each vertex to a layer according to the number of curves in A that pass above it; this is a breadth first layering starting from the vertex corresponding to the topmost (unbounded upward) connected component. No two vertices in the same layer have regions that project to overlapping subsets of the x-axis, so we may order the vertices within each layer according to the left-to-right ordering of these projections. This ordering is compatible with the planar embedding of the dual graph given by placing a representative point within each region and connecting each two adjacent regions by a curve crossing their shared boundary. □




Fig. 5. An arrangement of monotone curves whose dual graph has no planar layering

See Fig. 4 for examples of the graphs shown to have planar layerings by these theorems. Fig. 5 gives another example, demonstrating that Theorem 8 cannot be generalized to monotone curves whose projections do not cover the entire axis: it gives a family of monotone curves, all ending within the outer face of their arrangement, such that the dual graph of the arrangement has is not leveled planar. The dual graph is made of multiple $K_{2,3}$ subgraphs, each of which must have the 2-vertex side of its bipartition drawn on two layers with the 3-vertex side of its bipartition in a single layer between them; thus, up to top-bottom reflection, there is only a single layering for this graph that could possibly be planar. However, this layering forced by the planarity of the individual $K_{2,3}$ subgraphs is not planar globally, because it forces one of the two arms of the graph (upper and lower right) to collide with the "armpit" where the other arm meets the body of the graph (left). The graph is drawn without crossings in the figure, but in a way that does not respect any layering of the graph. This example is also a series-parallel graph, and shows that Theorem 6 cannot be generalized to series-parallel, treewidth-2, or 2-outerplanar graphs: none of these classes of graphs is leveled planar.

A.4 Outerplanar graphs

Theorem 9 (Felsner, Liotta, and Wismath [13]). *Every outerplanar graph has a weakly leveled planar drawing.*

Felsner et al. prove this result by a construction based on breadth-first search, using the BFS number and depth in the BFS tree as coordinates. Alternatively, Theorem 9 can be proven by using induction on the number of triangular faces of a maximal outerplanar graph to show that each such graph has a layout in which, on each edge of the outer face, there is room to add one more triangle with its new vertex one level below the upper level of the previous triangle vertices.




Fig. 6. A Halin graph drawn with all edges on a single level or consecutive levels

Felsner et al. used such a drawing, together with a spiral layout like the one in Fig. 1, to find an improper 3-track layout (allowing edges between consecutive vertices in a track) of any outerplanar graph. Dujmović et al. [9] proved that every outerplanar graph has a (proper) 5-track layout. Theorems 3 and 9 imply that every outerplanar graph has layered pathwidth at most 2.

A.5 Halin graphs

Recall that a *Halin graph* [28] is the graph formed from a tree with no degree-two vertices, embedded in the plane, by connecting the leaves of a tree by a cycle in the order given by the embedding. Di Giacomo and Meijer [26] proved that every Halin graph has a 5-track layout, and described a Halin graph with track-number at least 4. As far as we are aware, it is open whether every Halin graph has a 4-track layout.

Theorem 10. *Every Halin graph has a weakly leveled planar drawing.*

Proof. Choose an arbitrary leaf of the tree from which the Halin graph was constructed, to be the root of the tree. Then assign vertices to levels as follows: the root is assigned to level 0. Then, in stage i of the assignment ($i = 1, 2, \dots$) we assign to level i the previously-unassigned nodes that are either children of nodes at level $i - 1$, or that belong to a path from such a child to its leftmost or rightmost leaf descendant.

This level assignment (depicted in Fig. 6) is consistent with the given planar embedding of the tree and the Halin graph formed from the tree. Clearly, it embeds each two vertices that are adjacent in the tree to the same level or consecutive levels, because if one of the two adjacent vertices is assigned to level i in stage i but the other is not, then the second vertex will be one of the children of the first vertex assigned to level $i + 1$ in the next stage. Additionally, pairs of vertices that are adjacent in the outer cycle of the Halin graph must also belong to the same level or adjacent levels, because (with the exception of the two edges incident to the root, for which a similar argument is possible) each such pair of vertices must consist of the rightmost descendant of one child and




Fig. 7. Adding an apex to a complete binary tree produces a series-parallel graph with track-number 4 but arbitrarily large layered pathwidth.

the leftmost descendant of the next child of the lowest common ancestor of the two nodes. The two children of the common ancestor can only be one level apart, and the same follows for their extremal descendants. □

Theorems 3 and 10 imply that every Halin graph has layered pathwidth at most 2.

A.6 Series-parallel and tree-apex graphs

Although series-parallel graphs are in some sense intermediate in complexity between outerplanar graphs and Halin graphs (for instance, series-parallel graphs and outerplanar graphs have treewidth 2, whereas Halin graphs in general have treewidth 3), it is not true that every series-parallel graph has bounded layered pathwidth. In fact it is not true that every *tree-apex* graph has bounded layered pathwidth, that is, a graph that can be obtained from a tree by by adding a universal vertex that is adjacent to all other vertices (Fig. 7).

Theorem 11. *For every integer $h \geq 0$, there exists a series-parallel graph, in fact a tree-apex graph, that has a track-number at most* 4 *and layered pathwidth $\Theta(h)$.*

Proof. Consider the tree-apex graph G formed from a complete binary tree of height h by adding a universal vertex that is adjacent to all other vertices (Fig. 7). G is series-parallel, has track-number at most 4 (since the tree has a 3-track layout), and has pathwidth $\Theta(h)$. G does not have bounded layered pathwidth because the universal vertex forces every layering of this graph to use at most three layers (the one containing this vertex and at most two adjacent layers). Every path decomposition of G has a bag with $\Omega(h)$ vertices, and the largest of the three intersections of this bag with a layer must also have $\Omega(h)$ vertices. Therefore, G has layered pathwidth $\Omega(h)$. □




Fig. 8. The 6-boundaried graphs U_p (left) and L_q (right) from the proof of Theorem 13.

B Proof that leveled planarity is not finite state

Following Downey and Fellows [27], we define a *t-boundaried graph* to be a graph G with t designated *boundary* vertices labeled $1, 2, \ldots, t$. Given two t-boundaried graphs G_1 and G_2 we define their *gluing* $G_1 \oplus G_2$ by identifying each boundary vertex of G_1 with the boundary vertex of G_2 having the same label.

An n-ary *t-boundaried operator* $\otimes$ consists of a t-boundaried graph $G_\otimes = (V_\otimes, E_\otimes)$ and injections $f_i : \{1, \ldots, t\} \to V_\otimes$ for $1 \leq i \leq n$. Then for t-boundaried graphs $G_1, \ldots, G_n$ we define the t-boundaried graph $G_1 \otimes \cdots \otimes G_n$ by gluing each G_i to $G_\otimes$ after applying f_i to the boundary labels of $G_\otimes$. After the gluing the labels of G_i are forgotten.

It can be shown that there exists a *standard set* of t-boundaried operators on t-boundaried graphs that can be used to generate the set of all graphs of treewidth t. Furthermore, it is possible to convert (in linear time) a tree decomposition of width t into a parse tree that uses these standard operators; see Theorem 12.7.1 in [27]. Define $\mathcal{U}_t^{\text{small}}$ to be the *small universe* of t-boundaried graphs obtained by parse trees, using these standard operators. Given a family of graphs F, we define the equivalence relation $\sim_F$ on $\mathcal{U}_t^{\text{small}}$, such that $G_1 \sim_F G_2$ if and only if for all $H \in \mathcal{U}_t^{\text{small}}$, we have $G_1 \oplus H \in F \Leftrightarrow G_2 \oplus H \in F$.

A family of graphs F is said to be *t-finite state* if the family of parse trees for graphs in $F_t = F \cap \mathcal{U}_t^{\text{small}}$ is t-finite state. Equivalently, such a family of parse trees may be recognized by a finite tree automaton. We can now state the analog of the Myhill–Nerode Theorem (characterizing recognizability of sets of strings by finite state machines) for treewidth t graphs in place of strings and finite tree automata in place of finite state machines.

Theorem 12 (Theorem 12.7.2 of [27]). *Let F be a family of graphs. Then F is t-finite state if and only if $\sim_F$ has finite index over $\mathcal{U}_t^{small}$.*

As we now show, leveled planarity is not t-finite state when t is sufficiently large.

Theorem 13. *For all $t \geq 6$, the family of leveled planar graphs is not t-finite state.*

Proof. Let F be the family of leveled planar graphs. It suffices to prove the theorem in the case when $t = 6$. Consider the 6-boundaried graphs U_p and L_q shown in Fig. 8, and observe that $U_p \oplus L_q$ is leveled planar if and only if $p = q$. So $U_p \sim_F U_\ell$ if and only if $p = \ell$, which implies that $\sim_{F_6}$ does not have finite index and that in turn F is not 6-finite state by Theorem 12. □

Theorem 13 implies that (when $t \geq 6$) the parse trees of leveled planar graphs with treewidth t are not recognizable by tree automata. Therefore automata-based methods such as Courcelle's Theorem cannot be used to show leveled planarity to be fixed-parameter tractable with respect to treewidth. In particular, leveled planarity cannot be expressed using the forms of monadic second-order graph logic to which Courcelle's Theorem applies.

Additional References

24. Zachary Abel, Erik D. Demaine, Martin L. Demaine, David Eppstein, Anna Lubiw, and Ryuhei Uehara, *Flat foldings of plane graphs with prescribed angles and edge lengths*, 22nd Int. Symp. Graph Drawing (GD 2014), LNCS, vol. 8871, Springer-Verlag, 2014, pp. 272–283, doi:10.1007/978-3-662-45803-7_23.
25. Hans-Jürgen Bandelt, Victor Chepoi, and David Eppstein, *Combinatorics and geometry of finite and infinite squaregraphs*, SIAM Journal on Discrete Mathematics **24** (2010), no. 4, 1399–1440, doi:10.1137/090760301, MR 2735930.
26. Emilio Di Giacomo and Henk Meijer, *Track drawings of graphs with constant queue number*, Proc. 11th International Symp. on Graph Drawing (GD '03), LNCS, vol. 2912, Springer, 2004, pp. 214–225, doi:10.1007/978-3-540-24595-7_20, MR 2177595.
27. Rodney G. Downey and Michael R. Fellows, *Fundamentals of parameterized complexity*, Texts in Computer Science, Springer, 2013, doi:10.1007/978-1-4471-5559-1.
28. Rudolf Halin, *Studies on minimally n-connected graphs*, Combinatorial Mathematics and its Applications (Proc. Conf., Oxford, 1969) (London), Academic Press, 1971, pp. 129–136, MR 0278980.